

1. Introduction

The Baltic Sea supports the economy and receives the impact of some 85 million people in nine coastal nations. Discussion around Baltic environmental change has focused on two closely combined topics, i.e. climate change and eutrophication. According to IPCC the northern Europe will experience increased rainfall in near future (Alcamo *et al.* 2007) which increases leaching of nutrients (e.g. Justić *et al.* 2003, Graham 2004) and accelerates eutrophication of sea (BACC author team 2008). Although Baltic countries are financing extensive monitoring programmes, there are only few studies done between climate regulation, runoffs, nutrient loadings and nutrient enrichment into the seawater (BACC author team 2008).

We have earlier presented a chain-of-events between changes in the North Atlantic weather patterns and subsequent changes in the Baltic Sea runoff and salinity and, finally, in Baltic biota (e.g. Hänninen *et al.* 2000, Hänninen *et al.* 2003, Vuorinen *et al.* 2004). The same reasoning is here extended to general Baltic system regulation and actual nutrient enrichment processes in the sea. The study is divided into three main questions:

1. Will increasing the number of climate predictors in modelling improve our understanding of the Baltic Sea system regulation?
2. Is it possible to model Baltic nutrient loading only on the basis of runoff and nutrient concentrations in the incoming water masses?
3. Is it possible to model nutrient concentrations in seawater using loading as the predictor?

2. Materials and methods

We modelled various weather indices to describe climatic regulation effects. NAO (North Atlantic Oscillation) was defined as normalised air pressure difference between Azores and Iceland. AO (Arctic Oscillation) were 1000mb height anomalies poleward of 20°N. SLP described sea level air pressures in Iceland. Finally, we used wind speed data from Gotland's Hoburg (56°92'22N, 18°14'71E). The runoff data were monthly values (km³) of total freshwater discharges from the catchment area into the Baltic Sea divided according to HELCOM Baltic sub-drainage basins (Fig.1), and comprising both monitored river runoffs and estimates of non-monitored runoffs. Nutrient data were downloaded from MARE's Nest (<http://nest.su.se/dest>), which holds reviewed Baltic Sea monitoring records, gathered by surrounding Baltic countries, and stored as databank according to the protocols of HELCOM (Anonymous 2005). We averaged data sets over HELCOM sub-basins and pooled it into three vertical water layers: 0-21m, 21-70m and below 70m, representing the vertical environmental gradient typical for the Baltic Sea.

In the statistical analysis we applied Transfer Function models, also called dynamic regressions (Box & Jenkins 1976, Pankratz 1991), using the Scientific Computing Associates (SCA) software (Liu & Lattayak 2007). Transfer functions are able to relate the response of one series not only to its own past values, but also to the past and present values of the other related time series. In practice this is realized by merging the basic concepts of the general regression model with that of traditional ARIMA models. The analysis was conducted for the period 1970-2000.

3. Results and discussion

Resulting Transfer Function models fit well with the observed series (Tables 1&2). All the substantial parameters showed statistical significance, and coefficient of determination (r^2) values varied between 0.49 and 0.88, which are considered satisfactory in statistical time series analysis.

General runoff regulation

All indices had specific influence in the Baltic freshwater runoff regulation. The most obvious difference between models could be detected in observed time lags (Table 1A). Larger geographical area, in general, meant longer regulation effect, i.e. delayed response from weather effect to a response of runoff into the sea. In north-south direction, the northern areas showed lagged response indicating stronger winter effect in north and east. NAO was evidently the best index to explain general runoff regulation. AO showed inverse and much longer regulation response. Iceland SLP indicated weaker but very similar kind of regulation as NAO, but inversely. Also Hoburg's wind speeds resembled very much the NAO regulation.

Nutrient loading processes

Nutrient loading models indicated very strong coupling between nutrient loading and freshwater runoff (Table 1B). All models showed that loading had instant response to runoffs in every geographical area studied. NH₄-N showed decreasing trend during study period.

Nutrient concentrations in seawater

The effect of phosphorus loading can be seen in tot-P concentrations in upper and middle seawater layers (Fig.2), with a lag of about one year in the central Baltic Sea, and a bit longer in the Gulf of Bothnia (Table 2). None of our models for nitrogen manifested any connection between the nitrogen loading and concentrations in seawater, regardless of the chemical form (organic/inorganic) of the substance.

Table 2. Identified transfer function models for C) nutrient enrichment processes in seawater, initial estimates of the parameters with standard errors, t -values and p -values. For practicality, only significant models are presented. Coefficients of determination for the models are calculated with $r^2 = 1 - [(n-1)/(n-p)] \cdot (\text{sum of squares}_{\text{residual}} / \text{sum of squares}_{\text{total}})$, where n = number of observations and p = number of estimated parameters. The central Baltic Sea series were monthly series but quarterly in the Gulf of Bothnia. For more detailed description see Hänninen *et al.* 2000.

| C. Nutrient concentration models | | | | |
|----------------------------------------------------------------------------------------------------------------------------------|-------|--------|--------|--------|
| I. Gulf of Bothnia | | | | |
| Upper layer (0-21m) | | | | |
| $r^2 = 0.49, n = 113$ | | | | |
| tot-P concentration vs. tot-P loading | | | | |
| $(1-B^1) \text{tot-P}_{\text{conc},t} = (\omega_1 B^1) (1-B^1) \text{tot-P}_{\text{load},t} + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 0.73 | 0.80 | | |
| S.E. | 0.28 | 0.06 | | |
| t -value | 2.64 | 13.72 | | |
| p -value | 0.009 | <0.001 | | |
| II. Gulf of Bothnia | | | | |
| Middle layer (21-70m) | | | | |
| $r^2 = 0.57, n = 111$ | | | | |
| tot-P concentration vs. tot-P loading | | | | |
| $(1-B^1) \text{tot-P}_{\text{conc},t} = (\omega_1 B^1) (1-B^1) \text{tot-P}_{\text{load},t} + (1-\phi_1 B^1) (1-\phi_2 B^1) a_t$ | | | | |
| Estimate | 0.73 | 0.80 | 0.37 | |
| S.E. | 0.28 | 0.06 | 0.09 | |
| t -value | 2.64 | 12.30 | 4.16 | |
| p -value | 0.009 | <0.001 | <0.001 | |
| III. central Baltic Sea | | | | |
| Upper layer (0-21m) | | | | |
| $r^2 = 0.82, n = 343$ | | | | |
| tot-P concentration vs. tot-P loading | | | | |
| $(1-B^1) \text{tot-P}_{\text{conc},t} = (\omega_1 B^1) (1-B^1) \text{tot-P}_{\text{load},t} + (1-\phi_1 B^1) (1-\phi_2 B^1) a_t$ | | | | |
| Estimate | 0.30 | 0.68 | 0.36 | 0.22 |
| S.E. | 0.13 | 0.04 | 0.06 | 0.06 |
| t -value | 2.26 | 15.77 | 6.42 | 3.98 |
| p -value | 0.024 | <0.001 | <0.001 | <0.001 |
| IV. central Baltic Sea | | | | |
| Middle layer (21-70m) | | | | |
| $r^2 = 0.66, n = 343$ | | | | |
| tot-P concentration vs. tot-P loading | | | | |
| $(1-B^1) \text{tot-P}_{\text{conc},t} = (\omega_1 B^1) (1-B^1) \text{tot-P}_{\text{load},t} + (1-\phi_1 B^1) (1-\phi_2 B^1) a_t$ | | | | |
| Estimate | 0.40 | 0.60 | 0.23 | 0.21 |
| S.E. | 0.20 | 0.05 | 0.06 | 0.05 |
| t -value | 2.40 | 13.04 | 4.21 | 3.84 |
| p -value | 0.017 | <0.001 | <0.001 | <0.001 |




Fig.1. The total Baltic Sea with catchment area (thick line) and used subdivisions in modelling exercises. BB = Bothnian Bay, BS = Bothnian Sea (hereby = Gulf of Bothnia) and NBP = northern Baltic Proper, WBP = western Baltic Proper, EBP = eastern Baltic Proper (hereby = central Baltic Sea).

Table 1. Identified transfer function models for A) general runoff regulation by climate and B) riverine nutrient loading processes. Models indicate initial estimates of the parameters with standard errors and t -values with p -values. Coefficients of determination for the models are calculated with $r^2 = 1 - [(n-1)/(n-p)] \cdot (\text{sum of squares}_{\text{residual}} / \text{sum of squares}_{\text{total}})$, where n = number of observations and p = number of estimated parameters. All presented time series are monthly means. For more detailed description see Hänninen *et al.* 2000.

| A. Climate indices models | | | | |
|---------------------------------------------------------------------------------------------------------|---------|--------|--------|--------|
| I. total Baltic Sea | | | | |
| Runoff vs. NAO | | | | |
| $r^2 = 0.71, n = 359$ | | | | |
| $(1-B^1) \text{Runoff}_t = (\omega_1 + \omega_2 B^1) (1-B^1) \text{NAO}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 1.10 | 0.76 | 0.83 | 0.61 |
| S.E. | 0.18 | 0.18 | 0.03 | 0.04 |
| t -value | 6.00 | 4.19 | 25.88 | 14.38 |
| p -value | <0.001 | <0.001 | <0.001 | <0.001 |
| II. total Baltic Sea | | | | |
| Runoff vs. AO | | | | |
| $r^2 = 0.68, n = 359$ | | | | |
| $(1-B^1) \text{Runoff}_t = (\omega_1 B^1) (1-B^1) \text{AO}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | -0.55 | 0.81 | 0.60 | |
| S.E. | 0.26 | 0.03 | 0.04 | |
| t -value | -2.12 | 25.20 | 13.88 | |
| p -value | 0.035 | <0.001 | <0.001 | |
| III. total Baltic Sea | | | | |
| Runoff vs. Iceland SLP | | | | |
| $r^2 = 0.69, n = 359$ | | | | |
| $(1-B^1) \text{Runoff}_t = (\omega_1 + \omega_2 B^1) (1-B^1) \text{Iceland SLP}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | -0.17 | -0.15 | 0.81 | 0.62 |
| S.E. | 0.05 | 0.05 | 0.03 | 0.04 |
| t -value | -3.45 | -3.02 | 25.01 | 14.40 |
| p -value | <0.001 | 0.003 | <0.001 | <0.001 |
| IV. total Baltic Sea | | | | |
| Runoff vs. Hoburg Winds | | | | |
| $r^2 = 0.70, n = 359$ | | | | |
| $(1-B^1) \text{Runoff}_t = (\omega_1 + \omega_2 B^1) (1-B^1) \text{Wind}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 1.64 | 1.86 | 0.78 | 0.64 |
| S.E. | 0.38 | 0.37 | 0.03 | 0.04 |
| t -value | 4.37 | 4.97 | 22.96 | 15.35 |
| p -value | <0.001 | <0.001 | <0.001 | <0.001 |
| B. Nutrient loading models | | | | |
| I. total Baltic Sea | | | | |
| tot-N loading vs. Runoff | | | | |
| $r^2 = 0.88, n = 359$ | | | | |
| $(1-B^1) \text{tot-N}_t = \omega_1 (1-B^1) \text{Runoff}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 1614.46 | 0.81 | 0.50 | |
| S.E. | 71.86 | 0.03 | 0.05 | |
| t -value | 22.47 | 26.77 | 10.38 | |
| p -value | <0.001 | <0.001 | <0.001 | |
| II. total Baltic Sea | | | | |
| NH ₄ -N loading vs. Runoff | | | | |
| $r^2 = 0.82, n = 359$ | | | | |
| $(1-B^1) \text{NH}_4\text{-N}_t = -C + \omega_1 (1-B^1) \text{Runoff}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | -208.09 | 108.81 | 0.76 | 0.50 |
| S.E. | 41.11 | 11.10 | 0.04 | 0.05 |
| t -value | -5.06 | 9.81 | 21.51 | 10.55 |
| p -value | <0.001 | <0.001 | <0.001 | <0.001 |
| III. total Baltic Sea | | | | |
| NO ₃ -N loading vs. Runoff | | | | |
| $r^2 = 0.82, n = 359$ | | | | |
| $(1-B^1) \text{NO}_3\text{-N}_t = \omega_1 (1-B^1) \text{Runoff}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 977.26 | 0.75 | 0.37 | |
| S.E. | 56.87 | 0.04 | 0.05 | |
| t -value | 17.18 | 19.75 | 7.24 | |
| p -value | <0.001 | <0.001 | <0.001 | |
| IV. total Baltic Sea | | | | |
| tot-P loading vs. Runoff | | | | |
| $r^2 = 0.84, n = 359$ | | | | |
| $(1-B^1) \text{tot-P}_t = \omega_1 (1-B^1) \text{Runoff}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 69.15 | 0.85 | 0.55 | |
| S.E. | 1.47 | 0.03 | 0.05 | |
| t -value | 25.00 | 28.08 | 12.11 | |
| p -value | <0.001 | <0.001 | <0.001 | |
| IV. total Baltic Sea | | | | |
| PO ₄ -P loading vs. Runoff | | | | |
| $r^2 = 0.77, n = 359$ | | | | |
| $(1-B^1) \text{PO}_4\text{-P}_t = \omega_1 (1-B^1) \text{Runoff}_t + (1-\phi_1 B^1) a_t$ | | | | |
| Estimate | 34.07 | 0.87 | 0.44 | |
| S.E. | 1.47 | 0.03 | 0.05 | |
| t -value | 23.18 | 30.69 | 8.84 | |
| p -value | <0.001 | <0.001 | <0.001 | |




Fig.2. Models for tot-P concentrations in seawater on the basis of loadings. On each panel modeled (spots) and observed changes (open circles) of the time series are based on the identified TF-models. Letter next to topic refers to corresponding model in Table 2. Smooth red lines are drawn with distance-weighted least squares method. On small boxes of each panel are presented model fit scatterplots (observed values in X-axis, estimated values in Y-axis).

Acknowledgements

We wish to thank Miguel Rodriguez Medina/Stockholm University who contributed to this study at data compiling stage. For data we'd like to thank Per Stålnacke, SHMI, NOAA, Baltex, Estonian Environment Information Centre, Finnish Environment Institute, Southwest Finland Regional Environment Centre, Inst. für Miljöanalys/Sweden, Institute of Aquatic Ecology/Univ. of Latvia/Riga, Institute of Meteorology and Water Management/Gdynia/Poland, Joint Research Centre/Vilnius/Lithuania, MARE project, NW Administration of Roshydromet/Russia, Sea Fisheries Institute Gdynia/Poland and SIBER project. This study is a contribution to the "Assessment and Modelling of Baltic Ecosystem Response (AMBER)" -project financed by EU BONUS+ programme and "Developing Chemometrics with the Tools of Information Sciences (CHESS)" -project financed by the Tekes (MASI programme 658/31/06).

References

- ALCAMO, J., MORENO, J.M., NOVÁKY, B., BINDI, M., COROBOV, R., DEVOY, R.J.N., GIANNAKOPOLOS, C., MARTIN, E., OLESEN, J.E. & SHIDENKO, A. 2007. Europe. Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. In: Parry, M.L., Canziani, O.F., Palutikof, J.P., van der Linden, P.J. & Hanson, C.E., Eds., Cambridge University Press, Cambridge, UK.
- ANONYMOUS 2005. Nest - a decision support system for management of eutrophication in the Baltic Sea. A user manual, version 3.
- BACC AUTHOR TEAM 2008. Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies XXII. Springer-Verlag, Germany.
- BOX, G.E.P., & G.M. JENKINS. 1976. Time Series Analysis: Forecasting and Control [revised edition]. Holden Day, San Francisco, USA.
- GRAHAM, P. 2004. Climate change effects on river flow to the Baltic Sea. *Ambio* 33:235-241.
- HÄNNINEN, J., VUORINEN, I. & P. HJELT. 2000. Climatic factors in the Atlantic control the oceanographic and ecological changes in the Baltic Sea. *Limnology and Oceanography* 45(3):703-710.
- HÄNNINEN, J., VUORINEN, I., & KORNILOVS, G. 2003. Atlantic climatic factors control decadal dynamics of a Baltic Sea Copepod, *Temora longicornis*. *Ecography*, 26:672-678.
- JUSTIĆ, D., TURNER, R. E. & RABALAIS, N.N. 2003. Climatic influences on riverine nitrate flux: Implications for coastal marine eutrophication and hypoxia. *Estuaries* 26:1-11.
- LIU, L.M., LATTYAK, W. J. 2007. New and enhanced capabilities in Release 8 of the SCA statistical system. Scientific Computing Associates@Corp, Illinois, USA.
- PANKRATZ, A. 1991. Forecasting with Dynamic Regression Models. Wiley, New York, USA.
- VUORINEN, HÄNNINEN, J. & KORNILOVS, G. 2003. Transfer-function modelling between environmental variation and mesozooplankton in the Baltic Sea. *Progress in Oceanography* 59:339-356.